

# COVID-19 Vaccine Timing—Routine Schedule



Age*	Vaccine	If unvaccinated:	If had monovalent doses give bivalent (B) doses:
<b>6 months– 4 years</b>	<b>Pfizer– Infant/Toddler Bivalent</b>	→ 3-8 weeks →  → ≥8 weeks →	<b>If 1 prior dose, then:</b> 3-8** weeks  ≥8 weeks <b>If 2-3 prior doses, then:</b> ≥8 weeks
<b>6 months– 5 years</b>	<b>Moderna– Bivalent</b>	→ 4-8 weeks** → Use blue cap vial, 0.25mL	<b>If 1 prior dose, then<sup>^</sup>:</b> 4-8** weeks  Blue cap, 0.25mL <b>If 2 prior doses then<sup>^</sup>:</b> ≥2 months  Pink cap, 0.2mL
<b>6 + years<sup>β</sup></b>	<b>Moderna– Bivalent</b>	→ ≥4 months → Use blue cap vial, 6-11 years: 0.25mL, 12+ years: 0.5mL	
<b>5–11 years<sup>†</sup></b>	<b>Pfizer– Pediatric Bivalent</b>		
<b>12+ years</b>	<b>Pfizer/ Adol/Adult Bivalent</b>	→ ≥4 months →	<b>If 1 or more prior doses (of any of the 4 brands), then<sup>†</sup>:</b> ≥2 months (Ages 65+ may receive 1 additional bivalent dose ≥4 months after 1st)
<b>12+ years</b>	<b>Novavax<sup>‡</sup> Monovalent</b>	→ 3-8 weeks** →  → ↑	
<b>18+ years</b>	<b>Janssen (J&amp;J)<sup>§</sup> Monovalent</b>	→ ↑	

\* See [CDC recommendations](#) for children transitioning from a younger to older age group

\*\* An [8-week interval](#) may be preferable for some people, especially for males 12-39 years.

β Please note the Moderna bivalent dosages differ for ages 6-11 (0.25 mL, 25 mcg) and 12+ (0.5 mL, 50 mcg) years.

^ Children 5 years of age who had 1 or more doses of Moderna monovalent vaccine may receive Moderna or Pfizer bivalent vaccine.

† Children 5 years of age who had 1 or more doses of Pfizer monovalent vaccine are only eligible to receive Pfizer bivalent vaccine.

‡ People 18+ years who have not received any booster doses and are unable or unwilling to receive a bivalent booster, may receive the [monovalent Novavax booster as a single booster dose](#) at least 6 months after their primary series.

View [Interim Clinical Considerations for Use of COVID-19 Vaccines](#) for details. Schedule is subject to change.

## Children and Adolescents (Ages 6 months–17 years)

Age	Vaccine	If unvaccinated* <sup>§</sup> :	If had prior monovalent doses give bivalent (B) doses* <sup>§</sup> :
<b>6 months–4 years</b>	<b>Pfizer Bivalent–Infant/Toddler</b>		<p><b>1</b> Monovalent → 3 w → <b>B</b> → ≥8 w → <b>B</b> → ≥2 m → <b>Optional Dose*</b></p> <p><b>2-3</b> Monovalent → ≥8 w → <b>B</b> → ≥2 m → <b>Optional Dose*</b></p>
<b>5–11 years</b>	<b>Pfizer Bivalent–Pediatric</b>		<p><b>1</b> Monovalent<sup>†</sup> → 3 w → <b>B</b> → ≥4 w → <b>B</b> → ≥2 m → <b>Optional Dose*</b></p> <p><b>2</b> Monovalent<sup>†</sup> → ≥4 w → <b>B</b> → ≥2 m → <b>Optional Dose*</b></p> <p><b>3</b> Monovalent<sup>†</sup> → ≥8 w → <b>B</b> → ≥2 m → <b>Optional Dose*</b></p> <p>Orange cap: 5-11 yrs, Gray cap: 12+ yrs</p>
<b>12+ years</b>	<b>Pfizer Bivalent–Adol/Adult</b>	Use orange cap for 5-11 years and gray cap for 12+ years.	
<b>6+ months</b>	<b>Moderna Bivalent</b>	<p><b>Use blue cap vial, 6 months-11 years: 0.25mL, 12+ years: 0.5mL</b></p>	<p><b>1</b> Monovalent<sup>^</sup> → 4 w → <b>B</b> → ≥4 w → <b>B</b> → ≥2 m → <b>Optional Dose*</b></p> <p><b>2</b> Monovalent<sup>^</sup> → ≥4 w → <b>B</b> → ≥2 m → <b>Optional Dose*</b></p> <p><b>3</b> Monovalent<sup>^</sup> → ≥8 w → <b>B</b> → ≥2 m → <b>Optional Dose*</b></p> <p>Blue cap: 6+ yrs, Pink (or Blue) cap: 6 m-5 yrs</p>
<b>12+ years</b>	<b>Novavax Monovalent</b>		<p><b>If 1 or 2 prior doses, then:</b></p> <p>≥8 weeks → <b>Bivalent (Moderna/Pfizer)</b> → ≥2 m → <b>Optional Dose* (Moderna/Pfizer)</b></p>

\* An **optional dose** may be given ≥2 months after the last dose. **Further doses** may be given at the healthcare provider’s discretion. See [Table 2](#) for vial and dosage.

§ Heterologous “mix-and-match” dosing is allowed for ages 6 years and older.

† Children 5 years of age who had 1 or more doses of Pfizer monovalent vaccine are only eligible to receive Pfizer bivalent vaccine.

^ Children 5 years of age who had 1 or more doses of Moderna monovalent vaccine may receive Moderna or Pfizer bivalent vaccine.

View [Interim Clinical Considerations for Use of COVID-19 Vaccines](#) for details. Schedule is subject to change.

## Adults (Ages 18 years and older)

Vaccine	If unvaccinated*§:	If had prior monovalent doses give bivalent (B) doses*§:																	
<b>Moderna– Bivalent</b> (blue cap/gray border)	<p><b>Use blue cap vial, 0.5mL (50mcg), for 12+ years.</b></p>	<table border="1"> <tr> <td>1 Monovalent</td> <td>4 w</td> <td>B</td> <td>≥4 w</td> <td>B</td> <td rowspan="3">≥2 m</td> <td rowspan="3">Optional Dose* (Moderna/Pfizer)</td> </tr> <tr> <td>2 Monovalent</td> <td></td> <td></td> <td>≥4 w</td> <td>B</td> </tr> <tr> <td>3 Monovalent</td> <td></td> <td></td> <td>≥8 w</td> <td>B</td> </tr> </table>	1 Monovalent	4 w	B	≥4 w	B	≥2 m	Optional Dose* (Moderna/Pfizer)	2 Monovalent			≥4 w	B	3 Monovalent			≥8 w	B
1 Monovalent	4 w	B	≥4 w	B	≥2 m	Optional Dose* (Moderna/Pfizer)													
2 Monovalent			≥4 w	B															
3 Monovalent			≥8 w	B															
<b>Pfizer– Bivalent Adol/Adult</b> (gray cap)		<table border="1"> <tr> <td>1 Monovalent</td> <td>3 w</td> <td>B</td> <td>≥4 w</td> <td>B</td> <td rowspan="3">≥2 m</td> <td rowspan="3">Optional Dose* (Moderna/Pfizer)</td> </tr> <tr> <td>2 Monovalent</td> <td></td> <td></td> <td>≥4 w</td> <td>B</td> </tr> <tr> <td>3 Monovalent</td> <td></td> <td></td> <td>≥8 w</td> <td>B</td> </tr> </table>	1 Monovalent	3 w	B	≥4 w	B	≥2 m	Optional Dose* (Moderna/Pfizer)	2 Monovalent			≥4 w	B	3 Monovalent			≥8 w	B
1 Monovalent	3 w	B	≥4 w	B	≥2 m	Optional Dose* (Moderna/Pfizer)													
2 Monovalent			≥4 w	B															
3 Monovalent			≥8 w	B															
<b>Novavax‡</b> Monovalent		If 1 or 2 prior doses, then: ≥8 weeks → <b>Bivalent</b> (Moderna/Pfizer) → ≥2 m → <b>Optional Dose*</b> (Moderna/Pfizer)																	
<b>Janssen (J&amp;J)</b> Monovalent		If 1 or 2 prior doses, then: ≥4 weeks → <b>Bivalent</b> (Moderna/Pfizer) → ≥2 m → <b>Optional Dose*</b> (Moderna/Pfizer)																	

\* An optional dose may be given ≥2 months after the last dose. Further doses may be given at the healthcare provider’s discretion. See [Table 2](#) for vial and dosage.

§ Heterologous “mix-and-match” dosing is allowed for ages 6 years and older.

‡ People 18+ years who have not received any booster doses and are unable or unwilling to receive a bivalent booster, may receive the [monovalent Novavax booster as a single booster dose](#) at least 6 months after their primary series.

View [Interim Clinical Considerations for Use of COVID-19 Vaccines](#) for details. Schedule is subject to change.